



Nancy Y. Kiang (611 - NASA-GISS), Antígona Segura, Giovanna Tinetti, Govindjee, Robert E. Blankenship, Martin Cohen, Janet Siefert, David Crisp, Victoria S. Meadows
nkang at giss.nasa.gov, <http://vpl.astro.washington.edu>

Project Objective

What plausible colors might future telescopes see for photosynthesis on exoplanets orbiting stars hotter or cooler than our Sun?

What kind of light spectrum will alien photosynthesizers see?

Recent Results

In papers published last year in the journal *Astrobiology*, we examined how photosynthetic pigments are adapted to solar radiation, and we simulated the light spectrum as would be seen by organisms living on exoplanets around F, K, quiescent M, and flaring M stars. We found:


- Green chlorophyll dominates on Earth because oxygen from photosynthesis itself shifted the Solar spectrum at the Earth's surface.
- Plants around F stars would likely absorb more in the blue.
- Plants around M stars could prefer the infrared and be anoxygenic.

Project Description


Photosynthesis on extrasolar planets will be adapted to a light spectrum from stars of other types than our Sun, filtered through a different atmosphere.

On Earth, we found these rules: pigments will preferentially absorb photons that are either the most abundant, the shortest available wavelength (most energetic), or longest available wavelength (where the photosynthetic reaction center operates). The available light spectrum is itself influenced by atmospheric inputs from photosynthesis: O₂ and O₃ shift solar radiation to have most photon in the red at the Earth's surface -- right where the oxygenic photosynthesis reaction center is.

We calculated the surface incident photon flux spectrum on Earth-like planets orbiting these star types: F, K, a quiescent M dwarf, and a UV flaring M star. Atmospheric composition, assuming Earth-like surface fluxes, with and without oxygen, was calculated with a photochemical model (Pavlov and Kasting, 2002) coupled to a radiative-convective model (Pavlov, et al., 2000). Incident light spectra were calculated with the high-resolution radiative transfer model, SMART (Meadows & Crisp, 1996; Crisp, 1997).



Land plants often get too much light, so they need only chlorophyll and carotenoid pigments to harvest mostly blue and red photons. The ozone Chappuis band in the visible region shifts the solar incident photon flux density to have peak photons in the red. Underwater, algae and cyanobacteria have extra pigments, phycobilins, to pick up more green and yellow photons. In deep, dark, anoxic waters, anoxygenic bacteria harvest leftover near-infrared photons and what little visible photons they can get.



Oxygenic photosynthesis makes the light spectrum on F2V planets bluer, due to shifting by the ozone Chappuis band. Plants will have so much blue light, they might need sunscreen like anthocyanin, but blue-colored. Life starting out on planets around young UV-flaring M stars will need to keep under water for protection. When the star matures and flares die down, no ozone shield will be needed on land. The dominant photosynthesis could be anoxygenic in the infrared.

Conclusions

- This work will help to interpret the data from NASA's Terrestrial Planet Finder and the European Space Agency's (ESA) Darwin missions for signs of photosynthetic life.
- Why are plants green? We offer a new explanation based on the coevolution of Earth photosynthesis with the atmosphere: oxygen from early oxygenic photosynthesis in the oceans gave green plants the competitive advantage on land. On planets around M stars, the competitive advantage could be in the infrared.

Publications and References

- Crisp, D. (1997) Absorption of sunlight by water vapor in cloudy conditions: a partial explanation for the cloud absorption anomaly. *Geophys. Res. Lett.* 24(5), 571-574.
- Kiang, N.Y., J. Siefert, Govindjee, and R.E. Blankenship. (2007). "Spectral signatures of photosynthesis I: Review of Earth organisms," *Astrobiology Special Issue on M Stars*, 7(1): 222-251.
- Kiang, N.Y., A. Segura, G. Tinetti, Govindjee, R.E. Blankenship, M. Cohen, J. Siefert, D. Crisp, and V.S. Meadows. (2007). "Spectral signatures of photosynthesis II: coevolution with other stars and the atmosphere on extrasolar worlds," *Astrobiology Special Issue on M Stars*, 7(1): 252-274.
- Pavlov, A.A. and Kasting, J.F. (2002) Mass-independent fractionation of sulfur isotopes in Archean sediments: strong evidence for an anoxic Archean atmosphere. *Astrobiology* 2(1), 27-41.
- Pavlov, A.A., Kasting, J.F., Brown, L.L., Rages, K.A., and Freedman, R. (2000) Greenhouse warming by CH₄ in the atmosphere of early Earth. *J. Geophys. Res.* 105(E5), 11981-11990.
- Meadows, V.S., and Crisp, D. (1996) Ground-based nearinfrared observations of the Venus nightside: the thermal structure and water abundance near the surface. *J. Geophys. Res.* 101(E2), 4595-4622.